arXiv:1207.4540v1 [cond-mat.stat-mech] 19 Jul 2012

Finite size behaviors of critical Ising model on a rectangle with free boundaries

Xintian Wu,[1,*] Nickolay Izmailian,[1,2,†] and Wenan Guo[1,‡]

[1] *Department of Physics, Beijing Normal University, Beijing, 100875, China*
[2] *A.I. Alikhanyan National Science Laboratory, Alikhanian Br.2, 375036 Yerevan, Armenia.*

(Dated: July 20, 2012)

Using the bond-propagation algorithm, we study the Ising model on a rectangle of size $M \times N$ with free boundaries. For five aspect ratios $\rho = M/N = 1, 2, 4, 8, 16$, the critical free energy, internal energy and specific heat are calculated. The largest size reached is $M \times N = 64 \times 10^6$. The accuracy of the free energy reaches 10^{-26}. Basing on these accurate data, we determine exact expansions of the critical free energy, internal energy and specific heat. With these expansions, we extract the bulk, surface and corner parts of free energy, internal energy and specific heat. The fitted bulk free energy density is given by $f_\infty = 0.92965398834161021499(1)$, comparing with Onsager's exact result $f_\infty = 0.92965398834161021506\cdots$. We prove the conformal field theory(CFT) prediction of the corner free energy, in which the central charge of the Ising model is found to be $c = 0.5 \pm 1 \times 10^{-10}$ comparing with the CFT result $c = 0.5$. We find that not only the corner free energy but also the corner internal energy and specific heat are geometry independent, i.e., independent of aspect ratio. The implication of this finding on the finite scaling is discussed. In the second order correction of the free energy, we prove the geometry dependence predicted by CFT and find out a geometry independent constant beyond CFT. High order corrections are also obtained.

PACS numbers: 75.10.Nr,02.70.-c, 05.50.+q, 75.10.Hk

Introduction. Finite size effect has been attracting tremendous interest in the study of condensed matter physics. It becomes of practical interest due to the recent progresses in fine processing technologies, which has enabled the fabrication of nanoscale materials with novel shapes [1–3]. Exact solutions have been playing a key role in determining the form of finite size scaling. Ferdinand and Fisher [4] pioneered on the 2D Ising model on a finite size lattice, which extended Onsager's exact solution [5] and stimulated the ideas of finite size scaling. Since then, exact results of the model on finite size lattices with various boundaries have been studied intensively [4–12]. Detailed knowledge has been obtained for the torus case [8, 9], for helical boundary condition [10], for Brascamp-Kunz boundary condition [11, 12] and for infinitely long cylinder [13]. However in the jigsaw puzzle of the 2D Ising model an important piece is still missed, which is the exact result on a rectangle with free boundaries. Generally speaking the rectangle geometry system is very interesting in its own right. It is, for instance, the natural one to consider in the case of quenches for one dimensional quantum systems with open boundaries. In 2D it is the simplest geometry to study transport properties for Anderson localization. Although there are Monte Carlo and transfer matrix studies on this problem [14, 15], the accuracy or the system sizes of the results are not enough to extract the finite size corrections. Meanwhile, for 2D critical systems, a huge amount of knowledge has been obtained by the application of the powerful techniques of integrability and conformal field theory (CFT) [16–18]. Cardy and Peschel predicted that the next subdominant contribution to the free energy on a square comes from the corners [17], which is universal, and related to the central charge c in the continuum limit. Kleban and Vassileva [18] extended the study of the free energy on a rectangle. They further derived a geometry dependent term to the free energy. However, they did not determine a geometry independent additive constant in the coefficient. Till now there is few evidence for these predictions from exact solutions or numerical calculations. Furthermore, as far as we know, there is no detailed study of the internal energy and the specific heat on a rectangle neither by CFT nor by exact solutions/numerical calculations.

Recently an efficient bond propagation (BP) algorithm was developed for computing the partition function of the Ising model in two dimensions, which is exact to machine precision and works for any planar network of Ising spins with arbitrary bond strengths [19, 20]. It is also much faster than Monte Carlo simulation, and costs quite moderate memory comparing with the transfer matrix method. Very large system size can be reached. The BP algorithm is thus a powerful tool to study the Ising model on a rectangle with free edges and corners. In this letter we apply the BP algorithm to study the Ising model on an $M \times N$ rectangle. We obtain finite size data of the critical free energy f, internal energy U and specific heat c. By fitting these data we find that the exact expansion of the critical free energy, internal energy and specific heat can be written in the following form

$$f = f_\infty + f_{surf}\frac{M+N}{S} + f_{corn}\frac{\ln S}{S} + \sum_{k=1}^{\infty}\frac{A_k}{S^{(k+1)/2}}, \quad (1)$$

$$U = U_\infty + U_{surf}\frac{M\ln N + N\ln M}{S} + U_{corn}\frac{\ln S}{S} + \sum_{k=1}^{\infty}\frac{B_k}{S^{k/2}}, \quad (2)$$

$$c = A_0 \ln N + c_0 + c_{surf}\frac{M\ln N + N\ln M}{S} + c_{corn}\frac{\ln S}{S} + \sum_{k=1}^{\infty}\frac{D_k}{S^{k/2}}, \quad (3)$$

where $S = M \times N$ is the area of the system. f_∞, U_∞ are the bulk term, f_{surf}, U_{surf} the surface coefficient, f_{corn}, U_{corn} the corner coefficient for the free energy, internal energy respectively. c_{surf}, c_{corn} are the corresponding coefficients for the specific heat. We find the fitted values $f_\infty, f_{surf}, U_\infty, U_{surf}, A_0$ are excellently consistent to the exactly known results [4–8, 10, 11]. The corner free energy $f_{corn} = c/8$ [17, 18] is proved. More over the geometry independent constant, which is ignored in Kleban and Vassileva work [18], is determined. We also find the corner contributions U_{corn} and c_{corn}, which are independent of aspect ratio. As far as we know, no previous studies predict such terms. We start by introducing the BP algorithm briefly. Then we present our numerical results and analysis.

Method. The schematic of the BP algorithm is shown in Fig. 1. In this algorithm BP series reduction, BP Y-Δ transformation and its inverse are the building blocks. By successively integrating in and then integrating out spin degrees of freedom in a way that only introduces local changes to the network, this algorithm progressively moves degrees of freedom to an open edge of the network, where they are eliminated. The transformation in each step is exact. The numerical accuracy is limited by machine's precision, which is the round-off error 10^{-32} in the quadruple precision. The BP algorithm needs about N^3 steps to calculate the free energy of an $N \times N$ lattice (much faster than other numerical method). Therefore the total error is approximately $N^{3/2} \times 10^{-32}$. This estimation has been verified in the following way: We compared the results obtained using double precision, in which there are 16 effective decimal digits, and those using quadruple precision. Because the latter results are much more accurate than the formal, we can estimate the error in double precision results by taking the quadruple results as the exact results. We thus found that the error is about $N^{3/2} \times 10^{-16}$. In our calculation, the largest size reached is $M = N = 8000$, the round-off error is less than 10^{-26}. The partition function of the Ising model on a 2D square lattice is

$$Z = \sum_{\{\sigma_i\}} \exp\left(\beta \sum_{<i,j>} \sigma_i\sigma_j\right), \quad (4)$$

where the nearest neighbor couplings are dimensionless and β is the inverse temperature. The free energy density, internal energy per spin and specific heat density are defined by

$$f = \frac{\ln Z}{MN}, \quad U = \frac{\partial f}{\partial \beta}, \quad c = \beta^2\frac{\partial^2 f}{\partial \beta^2}, \quad (5)$$



FIG. 1. (Color online) (a) and (b) are building blocks of the BP algorithm: BP series and BP $\Delta - Y$ transformation, respectively. (c) The schematic of BP algorithm. (c1) The BP series is applied to the three spins and two bonds in red. (c2) The $\Delta - Y$ transformation is applied to the three spins and three bonds in red. (c6) Finally only two spins and one bond are left.

respectively. With the BP algorithm, we get the free energy density f at the critical $\beta_c = \frac{1}{2}\ln(1+\sqrt{2}) = 0.44068679\cdots$ directly. The internal energy and specific heat are calculated by using a differentiation method

$$U \approx -\frac{f(\beta_c+\Delta\beta) - f(\beta_c - \Delta\beta)}{2\Delta\beta},$$
$$c \approx \beta_c^2\frac{f(\beta_c+\Delta\beta) + f(\beta_c-\Delta\beta) - 2f(\beta_c)}{(\Delta\beta)^2}. \quad (6)$$

In our calculation, $\Delta\beta = 10^{-7}$ is used. The error in the calculation of U due to the finite $\Delta\beta$ is approximately $\frac{1}{3}(\Delta\beta)^2\partial^3 f/\partial\beta^3$, which is less than 10^{-11} for the lattice with $N \le 2000$. The error in the calculation of c due to finite $\Delta\beta$ is $\frac{1}{12}(\Delta\beta)^2\partial^4 f/\partial\beta^4$, which is less than 10^{-9} for $N \le 2000$. Another source of error is the accumulated error due to round-off, which is less than 10^{-26}. This error is amplified by $1/\Delta\beta(1/\Delta\beta^2)$ times in the calculations of U (c), which is thus around 10^{-19} (10^{-12}). Therefore the errors of U, c are mainly caused by the finite $\Delta\beta$ in the differentiation. In other words the accuracies of the free energy, internal energy and specific heat are $10^{-26}, 10^{-11}, 10^{-9}$ respectively.

The calculations have been carried out for various aspect ratios $\rho = M/N = 1, 2, 4, 8, 16$ on an $M \times N$ rectangular lattice. For $M = N$, the calculation was carried out from $N = 30$ to $N = 8000$. For $\rho = 2, 4, 8, 16$, the calculated lattice sizes are from $N = 30$ to $N = 2000$.

Critical free energy. We fit the data of free energy density with the formula given by Eq. (1) with k from 1 to 8. The fitting method is the Levenberg-Marquardt method [21] for nonlinear fit. The standard deviation (SD) is defined by $SD = \sqrt{\sum_i (f_i - f_i^{(fit)})^2/(n_d - n_f)}$

with f_i the numerical data, $f_i^{(fit)}$ the value given by the fitting formula, n_d the data number and n_f the number of fitting parameters. For all ρ, SD reaches 10^{-20}.

The high accuracy can be seen from the bulk value f_∞. For $\rho = 1$, the fitted values of f_∞ is 0.92969539834161021499(1). The asymptotic bulk value of the free energy density should be the same as the exact result given by Onsager[5], i.e., $f_\infty = \ln\sqrt{2} + \frac{2}{\pi}G = 0.92969539834161021506\cdots$, where $G = 1 - \frac{1}{3^2} + \frac{1}{5^2} - \frac{1}{7^2} + \cdots$. Our estimation is consistent to it in the accuracy of 10^{-20}. For other ρ the consistency is the same.

According to finite size scaling, the surface correction term stems from free edges. This correction for the model on an infinitely long strip with two free edges has been found through exact solution to be $(D_1 - \frac{1}{2}\ln(1+\sqrt{2}))/N$, where $D_1 = \int_0^\pi \ln[1 + \sqrt{2}(1-\cos\theta)^{1/2}(3-\cos\theta)^{-1/2}] \approx 0.2589553765253$ [7]. Note the infinitely long strip can be considered as a rectangle with a aspect ratio $\rho \to \infty$. Thus the surface correction for a rectangle should be $f_{surf}(M+N)/S$ with $f_{surf} = D_1 - \frac{1}{2}\ln(1+\sqrt{2}) = -0.1817314169844187569\cdots$. Our fitting results coincide with this prediction precisely. For $\rho = 1, 2$, the fitted values $-f_{surf}$ are 0.18173141698441877(3), 0.1817314169844188(1) respectively, and for $\rho = 4, 8, 16$, they are all 0.1817314169844188(2). The correction term $f_{corn}\ln S/S$ stems from the corners of the rectangle. According to Cardy and Peschel [17], the four corners, at $M = N$, give rise to the term $\frac{c}{4}\ln(N)/N^2$, where $c = 1/2$ is the central charge. Kleban and Vassileva [18] extended the results to a rectangle with $M \neq N$ using CFT: The corner free energy is $c\ln(S)/(8S)$, which yields

$$f_{corn} = \frac{c}{8} = 0.0625\ . \tag{7}$$

In our fitting f_{corn} are found to be $0.0625 + (2\pm3)\times10^{-14}$, $0.0625 + (1\pm2)\times10^{-12}$,$0.0625 + (1\pm5)\times10^{-12}$,$0.0625 + (1\pm5)\times10^{-12}$,$0.0625 \pm 1\times10^{-11}$ for $\rho = 1, 2, 4, 8, 16$ respectively. All of them lead to the central charge $c = 0.5$ with the error less than 10^{-10}. There is a $1/N^2$ correction for the infinitely long strip [7, 16] with the coefficient $c\pi/24 \approx 0.0654498$. Kleban and Vassileva [18] proved that, for a finite rectangle, the correction is proportional to $1/S = 1/(M\times N)$ with the coefficient

$$A_1' = -\frac{c}{4}\ln[\eta(q)\eta(q')], \tag{8}$$

where $\eta(q) = q^{1/24}\prod_{n=1}^\infty(1-q^n)$ with $q = e^{-2\pi\rho}, q' = e^{-2\pi/\rho}$. At the limit $\rho \to \infty$, $A_1'\rho^{-1} \to c\pi/24$ which recovers the infinitely long strip result.

However, Kleban and Vassileva mentioned that, in their derivation, a possible geometry-independent additive constant was ignored [18]. In other words, the coefficient of $1/S$ should be

$$A_1 = A_1' + F_0, \tag{9}$$

where F_0 is the constant which contributes $F_0/S = F_0/(\rho N^2)$ to the free energy density, which tends to 0 in the infinitely long strip limit. For $\rho = 1, 2, 4, 8, 16$, the values of A_1' are 0.065918017562, 0.087578866954, 0.175155990232, 0.393633679243, 0.873910756056 respectively. By comparing the fitted A_1 (see Tab. I) and the theoretical A_1' , we obtained this constant

$$F_0 = -0.0049488147(2). \tag{10}$$

The other parameters $A_2, \cdots, A_8$ are fitted and listed in Tab. I. For the infinitely long strip, $A_2\rho^{-3/2}, A_3\rho^{-2}$ and $A_4\rho^{-5/2}$, at the limit $\rho \to \infty$, correspond to the coefficients of N^{-3}, N^{-4} and N^{-5} terms in the finite size free energy expansion respectively, which have been obtained as $-0.04616(2), 0.024(1), 0.69(6)$, by using numerical transfer matrix techniques [22]. Simple extrapolations of the fitted values of A_2, A_3, A_4 show that the magnitude of A_2, A_3 agree with the transfer matrix results, but A_4 does not. We have tried other forms of formula to fit the critical free energy data. For example, we added the terms $\ln S/S^{3/2}, \ln S/S^2$ in the fitting formula and found that the corresponding coefficients are extremely small (less than 10^{-7}). We concluded that the logarithmic correction only appears in the corner term $\ln S/S$. We note here that, in the asymptotic expansion of the free energy for the Ising model with periodic boundary conditions [8], with Braskamp-Kunz boundary conditions [11] and with helical boundary conditions [10], only integer powers of S appear.

Critical internal energy. We fit the data of critical internal energy with the formula given by Eq. (2) with k from 1 to 4. The bulk value U_∞ is known to be $\sqrt{2} \approx 1.41421356237$ [5]. Our fit of U_∞ is 1.4142135624 for the five aspect ratios. The leading correction should be caused by the edges. In the exact result of Au-Yang and Fisher [7] on the strip with two free edges, the edges' correction is given by $-\frac{2}{\pi}\ln N/N$. We conjecture that, on the rectangle with four free edges, the edge correction is given by $-\frac{2}{\pi}(M\ln N + N\ln M)/(MN)$ with the coefficient $U_{surf}(\rho) = -\frac{2}{\pi} = -0.636619773\ldots$. This conjecture is proved by the fitted U_{surf} for all ρ ($U_{surf} = -0.6366198...$), which agree with the predicted value in the accuracy 10^{-7}.

The term $B_1/S^{1/2}$ is in fact scaled as $1/N$. In the infinity long strip limit, the coefficient before $1/N$ is known as $\frac{2}{\pi}(\frac{7}{2}\ln 2 + \gamma - \frac{\pi}{4} - \ln\pi) \approx 0.683158$ [7]. As we can see in Tab. II, $B_1/\sqrt{\rho}$ indeed approaches this limit as ρ increases.

Following the convention in the critical free energy, we write the coefficient of $(\ln S)/S$ as U_{corn}. We found

$$U_{corn}(\rho) \approx -0.45025(3), \tag{11}$$

which is independent of aspect ratio ρ. Apparently, this contribution becomes zero in the limit of infinitely long strip, in which there is no corner. From this point, it is rational to call this term corner's correction.

TABLE I. The fitted parameters of Eq. (1) for the critical free energy.

parameter	$\rho = 1$	$\rho = 2$	$\rho = 4$	$\rho = 8$	$\rho = 16$
A_1	0.060969202833(3)	0.08263005222(2)	0.170207175502(4)	0.388684864512(8)	0.86896194132(2)
A_2	0.0883883476(2)	0.0595556310(1)	−0.1009034881(4)	−0.666287857(1)	−2.423248445(3)
A_3	−0.0175362651(2)	0.04067435(2)	0.41842401(7)	2.1854954(3)	9.76558(1)
A_4	−0.02405666(2)	−0.174428(2)	−1.290133(9)	−8.15503(5)	−48.556(3)
A_5	0.066893(8)	0.4351(1)	3.9657(8)	33.676(6)	277.21(5)
A_6	−0.16558(2)	−1.147(4)	−13.85(5)	−161.7(5)	−1857(6)
A_7	0.3851(4)	3.22(8)	53(1)	86(2)0	137(4)00
A_8	−0.703(3)	−7.6(8)	−17(2)0	−387(4)0	−8(1)0000

TABLE II. The fitted parameters of Eq. (2) for the critical internal energy per spin.

parameter	$\rho = 1$	$\rho = 2$	$\rho = 4$	$\rho = 8$	$\rho = 16$
U_{surf}	−0.63661981(1)	−0.63661983(3)	−0.63661985(4)	−0.63661987(5)	−0.63661988(5)
B_1	−0.1213621(2)	−0.2586182(5)	−0.6453965(8)	−1.266482(1)	−2.151535(2)
U_{corn}	−0.450170(3)	−0.45018(1)	−0.45022(2)	−0.45028(4)	−0.45042(9)
B_2	−0.98604(2)	−1.1278(9)	−1.7138(2)	−3.2016(4)	−6.4893(9)
B_3	−0.2969(2)	−0.101(1)	0.833(3)	3.943(8)	13.40(2)
B_4	−0.040(2)	−0.41(1)	−2.40(4)	−11.2(2)	−47.7(7)

The other parameters B_1, B_2, B_3, B_4 are fitted and listed in Tab. II. Again we have tried other forms of formula to fit the critical internal energy. The terms $\ln S/S^{3/2}, \ln S/S^2$ are excluded considering the coefficients are extremely small. Moreover the standard deviations of the fits with these terms are much larger than those without them.

Critical specific heat. The data of the critical specific heat are fitted using the formula given by Eq. (3) with k from 1 to 4. The leading term $A_0 \ln N$ is known from Onsager's exact result [5], which reads $A_0 = \frac{2}{\pi}[\ln(1+\sqrt{2})]^2 \approx 0.494538589$. Our fitting gives $A_0 \approx 0.49453858$. The other fitted parameters are shown in Tab. III. The coefficient c_0 increases with the aspect ratio. For the strip case it is known that $-c_0(\rho=\infty) = (\frac{7}{2}\ln 2 + \gamma - 14\xi(3)/\pi^2 - \frac{\pi}{4} - \ln\pi) \approx 0.3125538$ [7]. c_0 approaches this limit as $\rho \to \infty$ obviously, see Tab. III. However we have not obtained an analytical expression for the dependence of c_0 on ρ.

The term $(M \ln N + N \ln M)/S$ is the next order correction. Its coefficient c_{surf} is independent of ρ, and its average value over ρ is $c_{surf} = 0.524529(3)$. Note that this term is absent in the torus case [4] and not mentioned in the long strip case [7], but exists in the cylinder case with Brascamp-Kunz boundary conditions [11, 12].

The corner term c_{corn} also seems independent of aspect ratio ρ and its average value is about

$$c_{corn} = 0.368(1). \tag{12}$$

We have also tried many other forms of formula to fit the critical specific heat data. For example, we added the terms $(\ln S)^3/S, (\ln S)^2/S$ in the fitting formula and found that their coefficients are extremely small.

Conclusion. Using the BP algorithm, we studied the Ising model on a rectangle of size $M \times N$ with free boundaries. For five aspect ratios $\rho = 1, 2, 4, 8, 16$, the critical free energy, internal energy and specific heat were calculated. We brought numerical evidence that the correction term f_{corn} stems from the corners of the rectangle is indeed universal and is proportional to the central charge c. We also found that the terms U_{corn} and c_{corn} are independent from the aspect ratio ρ. In order to check whether or not the terms U_{corn} and c_{corn} are universal quantities, it is useful to extend our study to the Ising model on other types of lattices, e.g., the triangular, honeycomb, Kagome lattices. In such studies, the BP algorithm which is suitable for any planar network of Ising spins with arbitrary bond strengths [19, 20] is still a powerful tool. In addition, we can enhance the accuracy of the internal energy and specific heat by using an extended BP algorithm to calculate the internal energy without the using of differentiation[20]. Moreover the finite size effects of the correlation can be investigated by using the site propagation algorithm [23]. As we have shown, the sharp corners induce remarkable effects in critical region not only on the free energy, but also on the internal energy and the specific heat. It is expected that the sharp corners can induce remarkable effects on other properties of finite size systems in critical regime, for example, thermal conductivity, electric conductivity, etc. All of these effects should be observable in experiments.

TABLE III. The fitted parameters of Eq. (3) for the critical specific heat.

parameter	$\rho = 1$	$\rho = 2$	$\rho = 4$	$\rho = 8$	$\rho = 16$
c_0	−0.57078599(3)	−0.44276294(2)	−0.37766115(2)	−0.34510742(2)	−0.32883055(2)
c_{surf}	0.524516(2)	0.524533(2)	0.524531(2)	0.524530(3)	0.524529(3)
D_1	−0.34928(3)	−0.29524(1)	−0.17826(1)	−0.05986(2)	0.03117(2)
c_{corn}	0.3683(7)	0.3704(3)	0.3696(7)	0.368(1)	0.365(3)
D_2	1.144(5)	1.293(2)	1.914(5)	3.42(1)	6.67(3)
D_3	0.02(4)	−0.13(2)	−1.25(7)	−4.8(2)	−15.4(6)
D_4	0.8(3)	0.7(2)	3.6(8)	16(3)	7(1)0

acknowledgment This work is supported by the National Science Foundation of China (NSFC) under Grant No. 11175018.

* wuxt@bnu.edu.cn
† izmail@yerphi.am
‡ waguo@bnu.edu.cn

[1] S. Kawata, H.-B. Sun, T. Tanaka, K. Takeda, Nature **4**12, 697 (2001).
[2] V.F. Puntes, K.M. Krishnan, A.P. Alivisatos, Science **2**91, 2115 (2001).
[3] Y. Yin, R.M. Rioux, C.K. Erdonmez, S. Hughes, G.A. Somorjai, A.P. Alivisatos, Science **3**04, 711 (2004).
[4] A. E. Ferdinand and M. E. Fisher, Phys. Rev. **185**, 832 (1969)
[5] L. Onsager, Phys. Rev. **65**, 117 (1944).
[6] B. Kaufman, Phys. Rev. **76**, 1232 (1949).
[7] H. Au-Yang and M. E. Fisher, Phys. Rev. B **11**, 3469 (1975).
[8] E. V. Ivashkevich, N. Sh. Izmailian and Chin-Kun Hu, J. Phys. A **35** (2002) 5543.
[9] J. Salas, J. Phys. A: Math. Gen. 35 1833 (2002).
[10] N. Sh. Izmailian and Chin-Kun Hu, Phys. Rev. E. **76**, 041118 (2007).
[11] N. Sh. Izmailian, K. B. Oganesyan and Chin-Kun Hu, Phys. Rev. E **65** (2002) 056132.
[12] W. Janke and R. Kenna, Phys. Rev. B **65**, 064110 (2002).
[13] N. Sh. Izmailian and Chin-Kun Hu, Phys. Rev. Letts. **86**, 5160 (2001).
[14] D. P. Landau, Phys. Rev. B **13**, 2997 (1976).
[15] Borko Stošić, Sava Milošević and H. Eugene Stanley, Phys. Rev. B **,** 16, 11466 (1990).
[16] H. W. J. Blöte, J. L. Cardy, and M. P. Nightingale, Phys. Rev. Lett. **56**, 742 (1986).
[17] J. L. Cardy and I. Peschel, Nucl. Phys. B **300**, 377 (1988).
[18] P. Kleban and I. Vassileva, J. Phys. A: Math. & Gen. **24**, 3407 (1991).
[19] Y. L. Loh and E.W. Carlson, Phys. Rev. Lett. **97**, 227205 (2006).
[20] Y. L. Loh, E.W. Carlson, and M. Y. J. Tan, Phys. Rev. B **76**, 014404 (2007).
[21] Detailed description of the algorithm can be found in, e.g., W. H. Press, *et. al.*, *Numerical Recipes in Fortran 90*, (Cambridge University Press, 1997).
[22] S. L. A. de Queiroz, Phys. Rev. E **84**, 031107 (2011).
[23] X. T. Wu and J. Y. Zhao, Phys. Rev. B **80**, 104402 (2009).